












| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|--|----------------------------|----------------------------|------------------------------|------------------------------|---------------|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Unintentional Injuries and Violence | | | | | |
| Rarely or never wore a bicycle helmet (among students who rode a bicycle during the past 12 months) | 80.5 (77.9-82.9) | 79.8 (77.5-81.8) | | | * |
| Rarely or never wore a seat belt (when riding in a car driven by someone else) | 13.3 (10.9-16.1) | 6.6 (5.3-8.1) | * | | |
| Rarely or never wore a seat belt when driving (among students who drive a car) | 15.1 (12.2-18.6) | 7.4 (5.6-9.7) | * | | |
| Rode with a driver who had been drinking alcohol one or more times (in a vehicle during the past 30 days) | 24.5 (22.4-26.8) | 24.3 (21.9-26.9) | | | * |
| Drove when drinking alcohol one or more times (a car or other vehicle during the past 30 days) | 15.1 (12.9-17.6) | 9.6 (8.1-11.4) | * | | |
| Texted or e-mailed while driving (among students who drove a car during the past 30 days) | 55.7 (52.3-59.1) | 55.8 (52.0-59.6) | | | * |
| Talked on a cell phone while driving (among students who drove a car during the past 30 days) | 63.0 (59.8-66.0) | 59.7 (55.8-63.5) | | | * |
| Carried a weapon on at least one day (e.g., a gun, knife, or club during the past 30 days) | 38.5 (36.0-41.0) | 12.4 (11.1-13.8) | * | | |
| Carried a gun on at least one day (during the past 30 days) | 16.8 (15.0-18.8) | 3.7 (3.0-4.7) | * | | |
| Carried a weapon on school property on at least one day (e.g., a gun, knife, or club during the past 30 days) | 15.0 (13.3-16.9) | 4.5 (3.6-5.5) | * | | |
| Did not go to school because they felt unsafe at school or on their way to or from school on at least one day (during the past 30 days) | 6.5 (5.6-7.5) | 11.1 (9.5-12.8) | | * | |
| Threatened or injured with a weapon on school property one or more times (e.g., a gun, knife, or club during the past 12 months) | 7.6 (6.6-8.7) | 4.8 (3.9-5.9) | * | | |
| In a physical fight one or more times (during the past 12 months) | 29.6 (27-32.0) | 15.4 (13.5-17.5) | * | | |

| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|--|---------------------|---------------------|------------------------------|------------------------------|---------------|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Unintentional Injuries and Violence | | | | | |
| Injured in a physical fight one or more times (treated by a doctor or nurse, during the past 12 months) | 2.8 (2.3-3.4) | 1.6 (1.1-2.3) | ✱ | | |
| In a physical fight on school property one or more times (during the past 12 months) | 10.2 (9.0-11.6) | 4.1 (3.3-5.2) | ✱ | | |
| Bullied on school property (during the past 12 months) | 22.3 (20.4-24.3) | 30.5 (28.5-32.6) | | ✱ | |
| Ever been electronically bullied (e.g., e-mail, chat rooms, instant messaging Web sites, or texting during the past 12 months) | 10.6 (9.5-11.8) | 25.9 (23.8-28.2) | | ✱ | |
| Been the victim of teasing or name calling because someone thought they were gay, lesbian, or bisexual (during the past 12 months) | 13.0 (11.7-14.3) | 13.3 (11.7-15.1) | | | ✱ |
| Felt sad or hopeless (almost every day for 2 or more weeks in a row that they stopped doing some usual activities during the past 12 months) | 18.0 (16.4-19.7) | 35.4 (32.4-38.5) | | ✱ | |
| Seriously considered attempting suicide (during the past 12 months) | 12.7 (11.2-14.4) | 21.1 (19.2-23.2) | | ✱ | |
| Made a plan about how they would attempt suicide (during the past 12 months) | 11.0 (9.9-12.3) | 16.4 (14.7-18.2) | | ✱ | |
| Attempted suicide one or more times (during the past 12 months) | 6.4 (5.5-7.5) | 9.3 (7.8-11.1) | | ✱ | |
| Suicide attempt resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the past 12 months) | 2.7 (2.1-3.4) | 2.5 (1.9-3.2) | | | ✱ |
| Ever physically forced to have sexual intercourse (when they did not want to) | 5.7 (4.8-6.8) | 11.9 (10.5-13.6) | | ✱ | |
| Physically hurt on purpose by someone they were dating one or more times (among students who dated during the past 12 months) | 6.6 (5.3-8.2) | 11.0 (9.6-12.7) | | ✱ | |

| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|---|----------------------------|----------------------------|------------------------------|---|---|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Unintentional Injuries and Violence | | | | | |
| Been forced to do sexual things they did not want to one or more times (among dating students during the past 12 months) | 6.4 (4.8-8.5) | 15.6 (13.6-17.9) | |  | |
| Ever been choked by someone or tried to choke themselves on purpose such as with a belt, towel, or rope, for the feeling it caused | 7.3 (6.2-8.6) | 7.5 (6.3-8.8) | | |  |
| Tobacco Use | | | | | |
| Ever tried cigarette smoking (even one or two puffs) | 43.4 (40.4-46.5) | 38.6 (35.1-42.1) | | |  |
| Smoked a whole cigarette for the first time before age 13 years | 10.7 (9.3-12.3) | 8.6 (6.7-11.1) | | |  |
| Smoked cigarettes on at least one day (during the past 30 days) | 16.0 (14.1-18.2) | 14.4 (12.1-17.0) | | |  |
| Smoked cigarettes on 20 or more days (during the past 30 days) | 6.3 (5.1-7.7) | 4.6 (3.6-6.0) | | |  |
| Smoked more than 10 cigarettes per day (among students who currently smoked cigarettes, on the days they smoked during the past 30 days) | 7.5 (5.1-10.8) | 3.7 (2.0-7.0) | | |  |
| Smoked cigarettes on school property on at least one day (during the past 30 days) | 4.0 (3.2-5.0) | 3.4 (2.4-4.8) | | |  |
| Ever smoked at least one cigarette every day for 30 days | 9.6 (8.2-11.2) | 8.7 (7.3-10.3) | | |  |
| Tried to quit smoking cigarettes (among students who currently smoked cigarettes, during the past 12 months) | 50.4 (45.5-57.1) | 55.1 (47.6-62.4) | | |  |
| Usually obtained their own cigarettes by buying them in a store or gas station (among students who were age <18 years and who currently smoked cigarettes, during the past 30 days) | 11.8 (8.2-16.8) | 5.7 (3.2-10.1) | | |  |

| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|---|---------------------|---------------------|------------------------------|------------------------------|---------------|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Tobacco Use | | | | | |
| Used chewing tobacco, snuff, or dip on at least one day (during the past 30 days) | 21.6 (18.5-25.0) | 4.5 (3.3-6.1) | * | | |
| Used chewing tobacco, snuff, or dip on school property on at least one day (during the past 30 days) | 13.7 (11.2-16.4) | 2.4 (1.7-3.3) | * | | |
| Smoked cigars, cigarillos, or little cigars on at least one day (during the past 30 days) | 20.7 (18.4-23.3) | 10.4 (9.0-12.1) | * | | |
| Smoked cigarettes; smoked cigars, cigarillos, or little cigars; or used chewing tobacco, snuff, or dip on at least one day (during the past 30 days) | 33.0 (29.5-36.6) | 20.2 (17.7-23.0) | * | | |
| Alcohol and Other Drug Use | | | | | |
| Ever had at least one drink of alcohol on at least one day (during their life) | 73.1 (70.6-75.5) | 72.5 (70.1-74.7) | | | * |
| Drank alcohol for the first time before age 13 years | 22.9 (20.5-25.5) | 16.6 (14.5-18.8) | * | | |
| Had at least one drink of alcohol on at least one day (during the past 30 days) | 37.2 (34.3-40.1) | 36.9 (34.2-39.8) | | | * |
| Had five or more drinks of alcohol in a row within a couple of hours on at least one day (during the past 30 days) | 25.2 (22.9-27.6) | 21.6 (19.2-24.2) | | | * |
| Usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol during the past 30 days) | 30.0 (26.1-34.2) | 42.9 (39.3-46.6) | | * | |
| Had at least one drink of alcohol on school property on at least one day (during the past 30 days) | 6.9 (5.8-8.3) | 4.4 (3.6-5.5) | * | | |
| Ever used marijuana one or more times (during their life) | 38.4 (35.2-41.6) | 36.9 (33.2-40.9) | | | * |

| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|--|----------------------------|----------------------------|------------------------------|------------------------------|---------------|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Alcohol and Other Drug Use | | | | | |
| Tried marijuana for the first time before age 13 years | 9.0 (7.4-10.9) | 6.6 (5.9-8.8) | | | ✱ |
| Used marijuana one or more times (during the past 30 days) | 22.1 (19.7-24.7) | 19.9 (17.3-22.8) | | | ✱ |
| Used marijuana on school property one or more times (during the past 30 days) | 6.4 (5.0-8.2) | 4.0 (3.1-5.1) | | | ✱ |
| Ever used any form of cocaine one or more times (during their life) | 7.7 (6.6-9.0) | 4.9 (4.0-6.0) | ✱ | | |
| Ever sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times (during their life) | 9.4 (8.1-10.9) | 10.4 (8.6-12.5) | | | ✱ |
| Ever sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times (during the past 30 days) | 3.7 (3.0-4.5) | 2.8 (2.2-3.7) | | | ✱ |
| Ever used heroin one or more times (during their life) | 3.3 (2.6-4.2) | 1.7 (1.3-2.4) | ✱ | | |
| Ever used methamphetamines one or more times (during their life) | 4.4 (3.5-5.4) | 2.7 (2.0-3.5) | | | ✱ |
| Ever used ecstasy one or more times (during their life) | 9.4 (8.1-11.0) | 6.6 (5.5-8.0) | ✱ | | |
| Ever took steroid pills or shots without a doctor's prescription one or more times (during their life) | 3.7 (3.0-4.6) | 1.4 (1.0-2.0) | ✱ | | |
| Ever took prescription drugs one or more times without a doctor's prescription (such as Oxycontin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, during their life) | 16.8 (15.0-18.8) | 15.3 (13.7-17.2) | | | ✱ |
| Ever used a needle to inject any illegal drug into their body one or more times (during their life) | 3.0 (2.4-3.9) | 1.7 (1.2-2.4) | | | ✱ |

| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|---|---------------------|---------------------|------------------------------|------------------------------|---------------|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Alcohol and Other Drug Use | | | | | |
| Offered, sold, or given an illegal drug by someone on school property (during the past 12 months) | 24.9 (23.0-26.9) | 20.6 (18.7-22.6) | | | * |
| Sexual Behaviors | | | | | |
| Ever had sexual intercourse | 46.0 (42.4-49.6) | 46.0 (42.5-49.6) | | | * |
| Had sexual intercourse for the first time before age 13 years | 5.7 (4.4-7.3) | 2.8 (2.2-3.6) | * | | |
| Had sexual intercourse with four or more persons (during their life) | 15.3 (13.1-17.8) | 13.9 (11.8-16.2) | | | * |
| Had sexual intercourse with at least one person (during the past 3 months) | 32.8 (30.0-35.8) | 35.4 (32.5-38.4) | | | * |
| Drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active) | 23.5 (20.1-27.2) | 20.0 (17.5-22.8) | | | * |
| Used a condom during last sexual intercourse (among students who were currently sexually active) | 65.9 (61.6-69.9) | 57.2 (53.0-61.4) | * | | |
| Used birth control pills to prevent pregnancy before last sexual intercourse (among students who were sexually active) | 20.7 (17.1-24.7) | 29.0 (24.9-33.4) | | * | |
| Did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active) | 6.6 (5.0-8.7) | 8.6 (6.6-11.1) | | | * |
| Were ever taught in school about AIDS or HIV infection | 83.4 (80.5-86.0) | 84.2 (81.8-86.2) | | | * |

| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|--|----------------------------|----------------------------|------------------------------|------------------------------|---------------|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Dietary Behaviors | | | | | |
| Overweight (students who were $\geq 85^{\text{th}}$ percentile but $< 95^{\text{th}}$ percentile for body mass index) | 14.6 (13.1-16.1) | 11.2 (9.7-12.9) | ✱ | | |
| Obese (students who were $\geq 95^{\text{th}}$ percentile for body mass index) | 12.9 (11.5-14.6) | 5.6 (4.7-6.7) | ✱ | | |
| Described themselves as slightly or very overweight | 21.5 (19.6-23.6) | 34.3 (32.4-36.3) | | ✱ | |
| Were trying to lose weight | 25.7 (23.3-28.3) | 57.9 (55.7-60.1) | | ✱ | |
| Did not eat for 24 or more hours to lose weight or to keep from gaining weight (during the past 30 days) | 7.9 (6.5-9.6) | 15.7 (14.1-17.4) | | ✱ | |
| Took diet pills, powders or liquids to lose weight or to keep from gaining weight (without a doctor's advice, during the past 30 days) | 4.3 (3.5-5.2) | 6.0 (5.1-7.1) | | | ✱ |
| Vomited or took laxatives to lose weight or to keep from gaining weight (during the past 30 days) | 3.0 (2.2-3.9) | 6.0 (5.1-6.9) | | ✱ | |
| Did not drink 100% fruit juices (during the past seven days) | 19.7 (18.2-21.3) | 24.4 (22.6-26.2) | | ✱ | |
| Did not eat fruit (during the past seven days) | 10.2 (9.1-11.5) | 7.1 (6.0-8.5) | ✱ | | |
| Did not eat green salad (during the past seven days) | 35.1 (32.9-37.3) | 24.1 (22.3-26.0) | ✱ | | |
| Did not eat potatoes (excluding French fries, fried potatoes, or potato chips during the past seven days) | 23.6 (21.2-26.2) | 30 (27.8-32.3) | | ✱ | |
| Did not eat carrots (during the past seven days) | 40.7 (38.6-42.8) | 42.3 (39.6-45.0) | | | ✱ |

| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|--|----------------------------|----------------------------|------------------------------|------------------------------|---------------|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Dietary Behaviors | | | | | |
| Did not eat other vegetables (excluding green salad, potatoes, or carrots, during the past seven days) | 15.8 (14.1-17.7) | 11.4 (9.9-13.0) | ✱ | | |
| Did not eat fruits and vegetables 5 or more times per day (during the past seven days) | 81.0 (79.3-82.6) | 82.9 (80.8-84.8) | | | ✱ |
| Did not eat fruits or drink 100% fruit juices two or more times per day (during the past seven days) | 71.5 (69.5-73.4) | 72.0 (69.5-74.5) | | | ✱ |
| Did not eat fruit or drink 100% fruit juices (during the past seven days) | 4.6 (3.9-5.5) | 3.5 (2.8-4.5) | | | ✱ |
| Did not eat vegetables three or more times per day (during the past seven days) | 86.1 (84.7-87.5) | 87.5 (85.5-89.1) | | | ✱ |
| Did not eat vegetables (during the past seven days) | 4.8 (3.8-5.9) | 2.9 (2.2-3.7) | ✱ | | |
| Drank a can, bottle, or glass of soda or pop one or more times per day (during the past seven days) | 23.3 (21.0-25.8) | 12.7 (10.7-15.1) | ✱ | | |
| Did not drink a can, bottle or glass of soda or pop (during the past seven days) | 19.8 (17.9-21.8) | 33.1 (30.1-36.1) | | ✱ | |
| Drank a can, bottle, or glass of diet soda or pop one or more times per day (during the past seven days) | 5.6 (4.8-6.6) | 5.2 (4.0-6.7) | | | ✱ |
| Drank three or more glasses of milk per day (during the past seven days) | 18.4 (16.4-20.5) | 10.5 (9.3-11.8) | ✱ | | |
| Did not drink milk during the past seven days | 10.5 (9.0-12.3) | 17.4 (15.9-19.0) | | ✱ | |
| Ate breakfast on all of the past seven days | 41.9 (39.2-44.8) | 38.6 (36.1-41.3) | | | ✱ |

| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|--|---------------------|---------------------|------------------------------|------------------------------|---------------|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Dietary Behaviors | | | | | |
| Did not eat breakfast on any of the past seven days | 11.7 (10.1-13.5) | 11.6 (10.1-13.3) | | | |
| Most of the time or always went hungry (during the past 30 days) | 3.8 (3.0-4.8) | 2.7 (2.2-3.4) | | | ✱ |
| Physical Activity | | | | | |
| Physically active at least 60 minutes per day on five or more days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the past seven days) | 61.4 (59.0-63.8) | 47.7 (45.0-50.3) | ✱ | | |
| Did not participate in at least 60 minutes of physical activity on any day (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the past seven days) | 9.0 (7.9-10.3) | 12.3 (10.8-14.0) | | ✱ | |
| Watched TV three or more hours per day (on an average school day) | 24.6 (22.6-26.6) | 20.4 (18.1-22.8) | | | ✱ |
| Played video or computer games or used a computer for something that was not school work three or more hours per day (on an average school day) | 31.3 (29.4-33.3) | 28.0 (26.3-29.8) | | | ✱ |
| Did not attend physical education (PE) classes (in an average week when they were in school) | 40.5 (36.5-44.5) | 50.9 (46.2-55.5) | | ✱ | |
| Attended physical education (PE) classes daily (in an average week when they were in school) | 38.7 (34.7-42.9) | 30.8 (26.5-35.5) | | | ✱ |
| Played on sports teams (run by their school or community groups during the past 12 months) | 65.7 (63.2-68.2) | 61.4 (58.9-63.9) | | | ✱ |

| 2013 Montana Youth Risk Behavior Survey – Gender Comparison | | | | | |
|--|---------------------|---------------------|------------------------------|------------------------------|---------------|
| Question | Male | Female | Male More Likely Than Female | Female More Likely than Male | No Difference |
| Other | | | | | |
| Ever told by a doctor or nurse that they had asthma | 19.2 (17.8-20.7) | 20.5 (18.9-22.2) | | | ✱ |
| Used an indoor tanning device (such as a sunlamp, sunbed, or tanning booth during the past 12 months) | 7.2 (6.1-8.4) | 20.0 (17.4-22.8) | | ✱ | |
| Received help from a resource teacher, speech therapist, or other special education teacher (at school during the past 12 months) | 12.5 (11.0-14.2) | 12.1 (10.7-13.8) | | | ✱ |

MONTANA
YOUTH RISK BEHAVIOR SURVEY
WWW.OPI.MT.GOV/YRBS